

Wakimoto realization of Drinfeld current for the elliptic quantum algebra $U_{q,p}(\widehat{sl}_3)$

December 4, 2008

Takeo KOJIMA

*Department of Mathematics, College of Science and Technology, Nihon University,
Surugadai, Chiyoda-ku, Tokyo 101-0062, JAPAN*

Abstract

We study a free field realization of the elliptic quantum algebra $U_{q,p}(\widehat{sl}_3)$ for arbitrary level k . We give the free field realization of elliptic analogue of Drinfeld current associated with $U_{q,p}(\widehat{sl}_3)$ for arbitrary level k . In the limit $p \rightarrow 0, q \rightarrow 1$ our realization reproduces Wakimoto realization for the affine Lie algebra \widehat{sl}_3 .

PACS numbers : 02.20.Sv, 02.20.Uw, 02.30.Ik

1 Introduction

The elliptic quantum group has been proposed in papers [1, 2, 3, 4, 5]. There are two types of elliptic quantum groups, the vertex type $\mathcal{A}_{q,p}(\widehat{sl}_N)$ and the face type $\mathcal{B}_{q,\lambda}(g)$, where g is a Kac-Moody algebra associated with a symmetrizable Cartan matrix. The elliptic quantum groups have the structure of quasi-triangular quasi-Hopf algebras introduced by V.Drinfeld [6]. H.Konno [7] introduced the elliptic quantum algebra $U_{q,p}(\widehat{sl}_2)$ as an algebra of the screening currents of the extended deformed Virasoro algebra in terms of the fusion SOS model [8]. M.Jimbo, H.Konno, S.Odake, J.Shiraishi [9] continued to

study the elliptic quantum algebra $U_{q,p}(\widehat{sl_2})$. They constructed the elliptic analogue of Drinfeld currents and identified $U_{q,p}(\widehat{sl_2})$ with the tensor product of $\mathcal{B}_{q,\lambda}(\widehat{sl_2})$ and a Heisenberg algebra \mathcal{H} . The elliptic quantum group $\mathcal{B}_{q,\lambda}(\widehat{sl_2})$ is a quasi-Hopf algebra while the elliptic algebra $U_{q,p}(\widehat{sl_2})$ is not. The intertwining relation of the vertex operator of $\mathcal{B}_{q,\lambda}(\widehat{sl_2})$ is based on the quasi-Hopf structure of $\mathcal{B}_{q,\lambda}(\widehat{sl_2})$. By the above isomorphism $U_{q,p}(\widehat{sl_2}) \simeq \mathcal{B}_{q,\lambda}(\widehat{sl_2}) \otimes \mathcal{H}$, we can understand "intertwining relation" of the vertex operator for the elliptic algebra $U_{q,p}(\widehat{sl_2})$. Along the above scheme the elliptic analogue of Drinfeld current of $U_{q,p}(\widehat{sl_2})$ is extended to those of $U_{q,p}(g)$ for non-twisted affine Lie algebra g [9, 10]. In this paper we are interested in higher-rank generalization of level k free field realization of the elliptic quantum algebra. For the elliptic algebra $U_{q,p}(\widehat{sl_2})$, there exist two kind of free field realizations for arbitrary level k , the one is parafermion realization [7, 9], the other is Wakimoto realization [16]. In this paper we are interested in the higher-rank generalization of Wakimoto realization of $U_{q,p}(\widehat{sl_2})$. We construct level k free field realization of Drinfeld current associated with the elliptic algebra $U_{q,p}(\widehat{sl_3})$. This gives the first example of arbitrary level free field realization of the higher-rank elliptic algebra. This free field realization can be applied for construction of the integrals of motion for the elliptic algebra $U_{q,p}(\widehat{sl_3})$. For this purpose, see references [17, 18, 19].

The organization of this paper is as follows. In section 2 we set the notation and introduce bosons. In section 3 we review the level k free field realization of the quantum group $U_q(\widehat{sl_3})$ [15]. In section 4 we give the level k free field realization of the elliptic quantum algebra $U_{q,p}(\widehat{sl_3})$. In appendix we summarize the normal ordering of the basic operators.

2 Boson

The purpose of this section is to set up the basic notation and to introduce the boson. In this paper we fix three parameters $q, k, r \in \mathbb{C}$. Let us set $r^* = r - k$. We assume $k \neq 0, -3$ and $\text{Re}(r) > 0, \text{Re}(r^*) > 0$. We assume q is a generic with $|q| < 1, q \neq 0$. Let us set a pair of parameters p and p^* by

$$p = q^{2r}, \quad p^* = q^{2r^*}.$$

We use the standard symbol of q -integer $[n]$ by

$$[n] = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

Let us set the elliptic theta function $\Theta_p(z)$ by

$$\begin{aligned}\Theta_p(z) &= (z; p)_\infty (p/z; p)_\infty (p; p)_\infty, \\ (z; p)_\infty &= \prod_{n=0}^{\infty} (1 - p^n z).\end{aligned}$$

It is convenient to work with the additive notation. We use the parametrization

$$\begin{aligned}q &= e^{-\pi\sqrt{-1}/r\tau}, \\ p &= e^{-2\pi\sqrt{-1}/\tau}, \quad p^* = e^{-2\pi\sqrt{-1}/\tau^*}, \quad (r\tau = r^*\tau^*), \\ z &= q^{2u}.\end{aligned}$$

Let us set Jacobi elliptic theta function $[u]_r, [u]_{r^*}$ by

$$[u]_r = q^{\frac{u^2}{r}-u} \frac{\Theta_p(z)}{(p; p)_\infty^3}, \quad [u]_{r^*} = q^{\frac{u^2}{r^*}-u} \frac{\Theta_{p^*}(z)}{(p^*; p^*)_\infty^3}.$$

The function $[u]_r$ has a zero at $u = 0$, enjoys the quasi-periodicity property

$$[u + r]_r = -[u]_r, \quad [u + r\tau]_r = -e^{-\pi\sqrt{-1}\tau - \frac{2\pi\sqrt{-1}u}{r}} [u]_r.$$

Let us set the delta-function $\delta(z)$ as formal power series.

$$\delta(z) = \sum_{n \in \mathbb{Z}} z^n.$$

Following [15] we introduce free bosons $a_n^1, a_n^2, b_n^1, b_n^2, b_n^3, c_n^1, c_n^2, c_n^3, (n \in \mathbb{Z}_{\neq 0})$.

$$[a_n^i, a_m^j] = \frac{[(k+3)n][A_{i,j}n]}{n} \delta_{n+m,0}, \quad [p_a^i, q_a^j] = (k+3)A_{i,j}, \quad (i, j = 1, 2), \quad (2.1)$$

$$[b_n^i, b_m^j] = -\frac{[n]^2}{n} \delta_{i,j} \delta_{n+m,0}, \quad [p_b^i, q_b^j] = -\delta_{i,j}, \quad (i, j = 1, 2, 3), \quad (2.2)$$

$$[c_n^i, c_m^j] = \frac{[n]^2}{n} \delta_{i,j} \delta_{n+m,0}, \quad [p_c^i, q_c^j] = \delta_{i,j}, \quad (i, j = 1, 2, 3). \quad (2.3)$$

Here we have used Cartan matrix $\begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$.

For parameters $a_1, a_2, b_1, b_2, b_3, c_1, c_2, c_3 \in \mathbb{R}$, we set the vacuum vector $|a, b, c\rangle$ of the Fock space $\mathcal{F}_{a_1 a_2 b_1 b_2 b_3 c_1 c_2 c_3}$ as following.

$$a_n^i |a, b, c\rangle = b_n^j |a, b, c\rangle = c_n^k |a, b, c\rangle = 0, \quad (i = 1, 2; j = 1, 2, 3), \quad (2.4)$$

$$\begin{aligned}
p_a^i |a, b, c\rangle &= a_i |a, b, c\rangle, \quad p_b^j |a, b, c\rangle = b_j |a, b, c\rangle, \quad p_c^j |a, b, c\rangle = c_j |a, b, c\rangle, \\
&\quad (i = 1, 2; j = 1, 2, 3; n > 0).
\end{aligned} \tag{2.5}$$

The Fock space $\mathcal{F}_{a_1 a_2 b_1 b_2 b_3 c_1 c_2 c_3}$ is generated by bosons $a_{-n}^1, a_{-n}^2, b_{-n}^1, b_{-n}^2, b_{-n}^3, c_{-n}^1, c_{-n}^2, c_{-n}^3$ for $n \in \mathbb{N}_{\neq 0}$. The dual Fock space $\mathcal{F}_{a_1 a_2 b_1 b_2 b_3 c_1 c_2 c_3}^*$ is defined in the same manner. In this paper we construct the elliptic analogue of Drinfeld current for $U_{q,p}(\widehat{sl_3})$ by these bosons a_n^i, b_n^j, c_n^j acting on the Fock space.

3 Free Field Realization of $U_q(\widehat{sl_3})$

The purpose of this section is to give the free field realization of the quantum affine algebra $U_q(\widehat{sl_3})$. We give a review of Wakimoto realization of $U_q(\widehat{sl_3})$ [15]. Let us set the bosonic operators $a_{\pm}^i(z), b_{\pm}^i(z), \gamma^i(z), \beta_s^i(z)$ by

$$a_{\pm}^i(z) = \pm(q - q^{-1}) \sum_{n>0} a_{\pm n}^i z^{\mp n} \pm p_a^i \log q, \quad (i = 1, 2), \tag{3.1}$$

$$b_{\pm}^i(z) = \pm(q - q^{-1}) \sum_{n>0} b_{\pm n}^i z^{\mp n} \pm p_b^i \log q, \quad (i = 1, 2, 3), \tag{3.2}$$

$$b^i(z) = - \sum_{n \neq 0} \frac{b_n^i}{[n]} z^{-n} + q_b^i + p_b^i \log z, \quad (i = 1, 2, 3), \tag{3.3}$$

$$c^i(z) = - \sum_{n \neq 0} \frac{c_n^i}{[n]} z^{-n} + q_c^i + p_c^i \log z, \quad (i = 1, 2, 3), \tag{3.4}$$

$$\gamma^i(z) = - \sum_{n \neq 0} \frac{(b+c)_n^i}{[n]} z^{-n} + (q_b^i + q_c^i) + (p_b^i + p_c^i) \log(-z), \quad (i = 1, 2, 3), \tag{3.5}$$

$$\beta_1^i(z) = b_+^i(z) - (b^i + c^i)(qz), \quad \beta_2^i(z) = b_-^i(z) - (b^i + c^i)(q^{-1}z), \quad (i = 1, 2, 3), \tag{3.6}$$

$$\beta_1^i(z) = b_+^i(z) + (b^i + c^i)(qz), \quad \beta_2^i(z) = b_-^i(z) + (b^i + c^i)(q^{-1}z), \quad (i = 1, 2, 3). \tag{3.7}$$

We give a free field realization of Drinfeld current for $U_q(\widehat{sl_3})$.

Definition 3.1 We define the bosonic operators $e_1^+(z), e_2^+(z), e_1^-(z), e_2^-(z)$ by

$$e_1^+(z) = \frac{-1}{(q - q^{-1})z} (e_1^{+,1}(z) - e_1^{+,2}(z)), \tag{3.8}$$

$$e_2^+(z) = \frac{-1}{(q - q^{-1})z} (e_2^{+,1}(z) - e_2^{+,2}(z) + e_2^{+,3}(z) - e_2^{+,4}(z)), \tag{3.9}$$

$$e_1^-(z) = \frac{-1}{(q - q^{-1})z} (e_1^{-,1}(z) - e_1^{-,2}(z) - e_1^{-,3}(z) + e_1^{-,4}(z)), \tag{3.10}$$

$$e_2^-(z) = \frac{-1}{(q - q^{-1})z} (e_2^{-,1}(z) - e_2^{-,2}(z) + e_2^{-,3}(z) - e_2^{-,4}(z)). \tag{3.11}$$

$$\psi_1^\pm(z) = : \exp \left(b_\pm^1(q^{\pm k}z) + b_\pm^1(q^{\pm(k+2)}z) + b_\pm^2(q^{\pm(k+3)}z) - b_\pm^3(q^{\pm(k+2)}z) + a_\pm^1(q^{\pm \frac{k+3}{2}}z) \right) :, \quad (3.12)$$

$$\psi_2^\pm(z) = : \exp \left(-b_\pm^1(q^{\pm(k+1)}z) + b_\pm^2(q^{\pm k}z) + b_\pm^3(q^{\pm(k+1)}z) + b_\pm^3(q^{\pm(k+3)}z) + a_\pm^2(q^{\pm \frac{k+3}{2}}z) \right) :, \quad (3.13)$$

Here we have set

$$e_1^{+,1}(z) = : \exp(\beta_1^1(z)) :, \quad (3.14)$$

$$e_1^{+,2}(z) = : \exp(\beta_2^1(z)) :, \quad (3.15)$$

$$e_2^{+,1}(z) = : \exp(\gamma^1(z) + \beta_1^2(z)) :, \quad (3.16)$$

$$e_2^{+,2}(z) = : \exp(\gamma^1(z) + \beta_2^2(z)) :, \quad (3.17)$$

$$e_2^{+,3}(z) = : \exp(\beta_1^3(qz) + b_+^2(z) - b_+^1(qz)) :, \quad (3.18)$$

$$e_2^{+,4}(z) = : \exp(\beta_2^3(qz) + b_+^2(z) - b_+^1(qz)) :, \quad (3.19)$$

$$e_1^{-,1}(z) = : \exp \left(\beta_4^1(q^{-k-2}z) + b_-^2(q^{-k-3}z) - b_-^3(q^{-k-2}z) + a_-^1(q^{-\frac{k+3}{2}}z) \right) :, \quad (3.20)$$

$$e_1^{-,2}(z) = : \exp \left(\beta_3^1(q^{k+2}z) + b_+^2(q^{k+3}z) - b_+^3(q^{k+2}z) + a_+^1(q^{\frac{k+3}{2}}z) \right) :, \quad (3.21)$$

$$e_1^{-,3}(z) = : \exp \left(\gamma^2(q^{k+2}z) + \beta_1^3(q^{k+2}z) + b_+^2(q^{k+3}z) - b_+^3(q^{k+2}z) + a_+^1(q^{\frac{k+3}{2}}z) \right) :, \quad (3.22)$$

$$e_1^{-,4}(z) = : \exp \left(\gamma^2(q^{k+2}z) + \beta_2^3(q^{k+2}z) + b_+^2(q^{k+3}z) - b_+^3(q^{k+2}z) + a_+^1(q^{\frac{k+3}{2}}z) \right) :, \quad (3.23)$$

$$e_2^{-,1}(z) = : \exp \left(\gamma^2(q^{-k-1}z) - \beta_3^1(q^{-k-1}z) + 2b_-^3(q^{-k-1}z) + a_-^2(q^{-\frac{k+3}{2}}z) \right) :, \quad (3.24)$$

$$e_2^{-,2}(z) = : \exp \left(\gamma^2(q^{-k-1}z) - \beta_4^1(q^{-k-1}z) + 2b_-^3(q^{-k-1}z) + a_-^2(q^{-\frac{k+3}{2}}z) \right) :, \quad (3.25)$$

$$e_2^{-,3}(z) = : \exp \left(\beta_4^3(q^{-k-3}z) + a_-^2(q^{-\frac{k+3}{2}}z) \right) :, \quad (3.26)$$

$$e_2^{-,4}(z) = : \exp \left(\beta_3^3(q^{k+3}z) + a_+^2(q^{\frac{k+3}{2}}z) \right) :. \quad (3.27)$$

Here the symbol $:\mathcal{O}:$ represents the normal ordering of \mathcal{O} . For example we have

$$: b_k b_l := \begin{cases} b_k^i b_l^i, & k < 0 \\ b_l^i b_k^i, & k > 0. \end{cases} \quad : p_b^i q_b^i := : q_b^i p_b^i := q_b^i p_b^i.$$

Theorem 3.1 [15] *The bosonic operators $e_i^\pm(z)$, $\psi_i^\pm(z)$, ($i = 1, 2$) satisfy the following commutation relations.*

$$(z_1 - q^{A_{i,j}} z_2) e_i^+(z_1) e_j^+(z_2) = (q^{A_{i,j}} z_1 - z_2) e_j^+(z_2) e_i^+(z_1), \quad (3.28)$$

$$(z_1 - q^{-A_{i,j}} z_2) e_i^-(z_1) e_j^-(z_2) = (q^{-A_{i,j}} z_1 - z_2) e_j^-(z_2) e_i^-(z_1), \quad (3.29)$$

$$[\psi_i^\pm(z_1), \psi_j^\pm(z_2)] = 0, \quad (3.30)$$

$$\begin{aligned} & (z_1 - q^{A_{i,j}-k} z_2)(z_1 - q^{-A_{i,j}+k} z_2) \psi_i^\pm(z_1) \psi_j^\mp(z_2) \\ &= (z_1 - q^{A_{i,j}+k} z_2)(z_1 - q^{-A_{i,j}-k} z_2) \psi_j^\mp(z_2) \psi_i^\pm(z_1), \end{aligned} \quad (3.31)$$

$$(z_1 - q^{\pm(A_{i,j}-\frac{k}{2})} z_2) \psi_i^\pm(z_1) e_j^\pm(z_2) = (q^{\pm A_{i,j}} z_1 - q^{\mp \frac{k}{2}} z_2) e_j^\pm(z_2) \psi_i^\pm(z_1), \quad (3.32)$$

$$(z_1 - q^{\pm(A_{i,j}-\frac{k}{2})} z_2) e_i^\pm(z_1) \psi_j^\mp(z_2) = (q^{\pm A_{i,j}} z_1 - q^{\mp \frac{k}{2}} z_2) \psi_j^\mp(z_2) e_i^\pm(z_1), \quad (3.33)$$

$$\begin{aligned} & \{e_i^\pm(z_1) e_i^\pm(z_2) e_j^\pm(z_3) - (q + q^{-1}) e_i^\pm(z_1) e_j^\pm(z_3) e_j^\pm(z_2) + e_i^\pm(z_3) e_i^\pm(z_1) e_j^\pm(z_2)\} \\ & + \{z_1 \leftrightarrow z_2\} = 0, \quad \text{for } (i \neq j), \end{aligned} \quad (3.34)$$

$$[e_i^+(z_1), e_j^-(z_2)] = \frac{\delta_{i,j}}{(q - q^{-1}) z_1 z_2} \left(\delta \left(q^{-k} \frac{z_1}{z_2} \right) \psi_i^+(q^{-\frac{k}{2}} z_1) - \delta \left(q^k \frac{z_1}{z_2} \right) \psi_i^-(q^{-\frac{k}{2}} z_2) \right). \quad (3.35)$$

Hence $e_i^\pm(z), \psi_i^\pm(z)$ give level k free field realization of $U_q(\widehat{sl_3})$.

4 Free Field Realization of $U_{q,p}(\widehat{sl_3})$

The purpose of this section is to give a free field realization of the elliptic analogue of Drinfeld current for $U_{q,p}(\widehat{sl_3})$ with arbitrary level $k \neq 0, -3$. Let us set the bosonic operators $\mathcal{B}_\pm^{*i}(z), \mathcal{B}_\pm^i(z), (i = 1, 2, 3), \mathcal{A}^{*i}(z), \mathcal{A}^i(z), (i = 1, 2)$ by

$$\mathcal{B}_\pm^{*i}(z) = \exp \left(\pm \sum_{n>0} \frac{b_{-n}^i}{[r^*n]} z^n \right), \quad (i = 1, 2, 3), \quad (4.1)$$

$$\mathcal{B}_\pm^i(z) = \exp \left(\pm \sum_{n>0} \frac{b_n^i}{[rn]} z^{-n} \right), \quad (i = 1, 2, 3), \quad (4.2)$$

$$\mathcal{A}^{*i}(z) = \exp \left(\sum_{n>0} \frac{a_{-n}^i}{[r^*n]} z^n \right), \quad (i = 1, 2), \quad (4.3)$$

$$\mathcal{A}^i(z) = \exp \left(- \sum_{n>0} \frac{a_n^i}{[rn]} z^{-n} \right), \quad (i = 1, 2). \quad (4.4)$$

Definition 4.1 *Let us set the bosonic operators $e_i(z), f_i(z), \Psi_i^\pm(z), (i = 1, 2)$ by*

$$e_i(z) = U^{*i}(z)e_i^+(z), \quad (i = 1, 2), \quad (4.5)$$

$$f_i(z) = e_i^-(z)U^i(z), \quad (i = 1, 2), \quad (4.6)$$

$$\Psi_i^+(z) = U^{*i}(q^{\frac{k}{2}}z)\psi_i^+(z)U^i(q^{-\frac{k}{2}}z), \quad (i = 1, 2), \quad (4.7)$$

$$\Psi_i^-(z) = U^{*i}(q^{-\frac{k}{2}}z)\psi_i^-(z)U^i(q^{\frac{k}{2}}z), \quad (i = 1, 2). \quad (4.8)$$

Here we have set

$$U^{*1}(z) = \mathcal{B}_+^{*1}(q^{r^*}z)\mathcal{B}_+^{*1}(q^{r^*-2}z)\mathcal{B}_+^{*2}(q^{r^*-3}z)\mathcal{B}_-^{*3}(q^{r^*-2}z)\mathcal{A}^{*1}(q^{r^*+\frac{k-3}{2}}z), \quad (4.9)$$

$$U^{*2}(z) = \mathcal{B}_+^{*3}(q^{r^*-3}z)\mathcal{B}_+^{*3}(q^{r^*-1}z)\mathcal{B}_+^{*2}(q^{r^*}z)\mathcal{B}_-^{*1}(q^{r^*-1}z)\mathcal{A}^{*2}(q^{r^*+\frac{k-3}{2}}z), \quad (4.10)$$

$$U^1(z) = \mathcal{B}_-^1(q^{-r^*}z)\mathcal{B}_-^1(q^{-r^*+2}z)\mathcal{B}_-^2(q^{-r^*+3}z)\mathcal{B}_+^3(q^{-r^*+2}z)\mathcal{A}^1(q^{-r^*-\frac{k-3}{2}}z), \quad (4.11)$$

$$U^2(z) = \mathcal{B}_-^3(q^{-r^*+1}z)\mathcal{B}_-^3(q^{-r^*+1}z)\mathcal{B}_-^2(q^{-r^*}z)\mathcal{B}_+^1(q^{-r^*+1}z)\mathcal{A}^2(q^{-r^*-\frac{k-3}{2}}z). \quad (4.12)$$

The above free field realization of the twistors $U^{*i}(z), U^i(z), (i = 1, 2)$ is the main result of this paper.

Proposition 4.1 *The bosonic operators $e_i(z), f_i(z), \Psi_i^\pm(z), (i = 1, 2)$ satisfy the following commutation relations.*

$$e_i(z_1)e_j(z_2) = q^{-A_{i,j}} \frac{\Theta_{p^*}(q^{A_{i,j}}z_1/z_2)}{\Theta_{p^*}(q^{-A_{i,j}}z_1/z_2)} e_j(z_2)e_i(z_1), \quad (4.13)$$

$$f_i(z_1)f_j(z_2) = q^{A_{i,j}} \frac{\Theta_p(q^{-A_{i,j}}z_1/z_2)}{\Theta_p(q^{A_{i,j}}z_1/z_2)} f_j(z_2)f_i(z_1), \quad (4.14)$$

$$\Psi_i^\pm(z_1)\Psi_j^\pm(z_2) = \frac{\Theta_p(q^{-A_{i,j}}z_1/z_2)\Theta_{p^*}(q^{A_{i,j}}z_1/z_2)}{\Theta_p(q^{A_{i,j}}z_1/z_2)\Theta_{p^*}(q^{-A_{i,j}}z_1/z_2)} \Psi_j^\pm(z_2)\Psi_i^\pm(z_1), \quad (4.15)$$

$$\Psi_i^\pm(z_1)\Psi_j^\mp(z_2) = \frac{\Theta_p(pq^{-A_{i,j}-k}z_1/z_2)\Theta_{p^*}(p^*q^{A_{i,j}+k}z_1/z_2)}{\Theta_p(pq^{A_{i,j}-k}z_1/z_2)\Theta_{p^*}(p^*q^{-A_{i,j}+k}z_1/z_2)} \Psi_j^\mp(z_2)\Psi_i^\pm(z_1), \quad (4.16)$$

$$\Psi_i^\pm(z_1)e_j(z_2) = \frac{\Theta_{p^*}(q^{A_{i,j}\pm\frac{k}{2}}z_1/z_2)}{\Theta_{p^*}(q^{-A_{i,j}\pm\frac{k}{2}}z_1/z_2)} e_j(z_2)\Psi_i^\pm(z_1), \quad (4.17)$$

$$\Psi_i^\pm(z_1)f_j(z_2) = \frac{\Theta_{p^*}(q^{-A_{i,j}\mp\frac{k}{2}}z_1/z_2)}{\Theta_{p^*}(q^{A_{i,j}\mp\frac{k}{2}}z_1/z_2)} e_j(z_2)\Psi_i^\pm(z_1), \quad (4.18)$$

$$[e_i(z_1), f_j(z_2)] = \frac{\delta_{i,j}}{(q - q^{-1})z_1z_2} \left(\delta \left(q^{-k} \frac{z_1}{z_2} \right) \Psi_i^+(q^{-k/2}z_1) - \delta \left(q^k \frac{z_1}{z_2} \right) \Psi_i^-(q^{-k/2}z_2) \right), \quad (i \neq j). \quad (4.19)$$

We introduce the Heisenberg algebra \mathcal{H} generated by the following P_i, Q_i , ($i = 1, 2$).

$$[P_i, Q_j] = \frac{A_{i,j}}{2}, \quad (i, j = 1, 2). \quad (4.20)$$

Definition 4.2 Let us define the bosonic operators $E_i(z), F_i(z), H_i^\pm(z) \in U_q(\widehat{sl_3}) \otimes \mathcal{H}$, ($i = 1, 2$) by

$$E_1(z) = e_1(z) e^{2Q_1} z^{-\frac{P_1-1}{r^*}}, \quad E_2(z) = e_2(z) e^{2Q_2} z^{-\frac{P_2-1}{r^*}}, \quad (4.21)$$

$$F_1(z) = f_1(z) z^{\frac{2p_b^1+p_b^2-p_b^3+p_a^1}{r}} z^{\frac{P_1-1}{r}}, \quad F_2(z) = f_2(z) z^{\frac{2p_b^3+p_b^2-p_b^1+p_a^2}{r}} z^{\frac{P_2-1}{r}}, \quad (4.22)$$

$$H_1^\pm(z) = \Psi_1^\pm(z) e^{2Q_1} (q^{\mp \frac{k}{2}} z)^{\frac{2p_b^1+p_b^2-p_b^3+p_a^1}{r}} (q^{\pm(r-\frac{k}{2})} z)^{\frac{P_1-1}{r} - \frac{P_1-1}{r^*}}, \quad (4.23)$$

$$H_2^\pm(z) = \Psi_2^\pm(z) e^{2Q_2} (q^{\mp \frac{k}{2}} z)^{\frac{2p_b^3+p_b^2-p_b^1+p_a^2}{r}} (q^{\pm(r-\frac{k}{2})} z)^{\frac{P_2-1}{r} - \frac{P_2-1}{r^*}}. \quad (4.24)$$

Theorem 4.2 The bosonic operators $E_i(z), F_i(z), H_i^\pm(z)$, ($i = 1, 2$) satisfy the following commutation relations.

$$E_i(z_1) E_j(z_2) = \frac{\left[u_1 - u_2 + \frac{A_{i,j}}{2} \right]}{\left[u_1 - u_2 - \frac{A_{i,j}}{2} \right]_{r^*}} E_j(z_2) E_i(z_1), \quad (4.25)$$

$$F_i(z_1) F_j(z_2) = \frac{\left[u_1 - u_2 - \frac{A_{i,j}}{2} \right]}{\left[u_1 - u_2 + \frac{A_{i,j}}{2} \right]_r} F_j(z_2) F_i(z_1), \quad (4.26)$$

$$H_i^\pm(z_1) H_j^\pm(z_2) = \frac{\left[u_1 - u_2 - \frac{A_{i,j}}{2} \right]_r \left[u_1 - u_2 + \frac{A_{i,j}}{2} \right]}{\left[u_1 - u_2 + \frac{A_{i,j}}{2} \right]_r \left[u_1 - u_2 - \frac{A_{i,j}}{2} \right]_{r^*}} H_j^\pm(z_2) H_i^\pm(z_1), \quad (4.27)$$

$$H_i^+(z_1) H_j^-(z_2) = \frac{\left[u_1 - u_2 - \frac{A_{i,j}}{2} - \frac{k}{2} \right]_r \left[u_1 - u_2 + \frac{A_{i,j}}{2} + \frac{k}{2} \right]}{\left[u_1 - u_2 + \frac{A_{i,j}}{2} - \frac{k}{2} \right]_r \left[u_1 - u_2 - \frac{A_{i,j}}{2} + \frac{k}{2} \right]_{r^*}} H_j^-(z_2) H_i^+(z_1), \quad (4.28)$$

$$H_i^\pm(z_1) E_j(z_2) = \frac{\left[u_1 - u_2 \pm \frac{k}{4} + \frac{A_{i,j}}{2} \right]}{\left[u_1 - u_2 \pm \frac{k}{4} - \frac{A_{i,j}}{2} \right]_{r^*}} E_j(z_2) H_i^\pm(z_1), \quad (4.29)$$

$$H_i^\pm(z_1) F_j(z_2) = \frac{\left[u_1 - u_2 \mp \frac{k}{4} - \frac{A_{i,j}}{2} \right]}{\left[u_1 - u_2 \mp \frac{k}{4} + \frac{A_{i,j}}{2} \right]_r} F_j(z_2) H_i^\pm(z_1), \quad (4.30)$$

$$[E_i(z_1), F_j(z_2)] = \frac{\delta_{i,j}}{(q - q^{-1})z_1 z_2} \left(\delta \left(q^{-k} \frac{z_1}{z_2} \right) H_i^+(q^{-\frac{k}{2}} z_1) - \delta \left(q^k \frac{z_1}{z_2} \right) H_i^-(q^{-\frac{k}{2}} z_2) \right) \quad (4.31)$$

Now we have constructed level k free field realization of Drinfeld current $E_i(z), F_i(z), H_i^\pm(z)$ for the elliptic algebra $U_{q,p}(\widehat{sl_3})$. This gives the first example of arbitrary-level free field realization of higher-rank elliptic algebra.

Acknowledgement

The author would like to thank the organizing committee of the 27-th International Colloquium of the Group Theoretical Method in Physics held at Yerevan, Armenia 2008. The author would like to thank Prof.A.Kluemper for his kindness at Armenia. This work is partly supported by the Grant-in Aid for Young Scientist **B**(18740092) from Japan Society for the Promotion of Science.

Appendix

In appendix we summarize the normal ordering of the basic operators.

$$\begin{aligned} : e^{\gamma^i(z_1)} : \mathcal{B}_+^{*i}(z_2) &= : e^{\gamma^i(z_1)} \mathcal{B}_+^{*i}(z_2) : \frac{(q^{r^*+1} z_2/z_1; p^*)_\infty}{(q^{r^*-1} z_2/z_1; p^*)_\infty}, \\ : e^{\beta_1^i(z_1)} : \mathcal{B}_+^{*i}(z_2) &= : e^{\beta_1^i(z_1)} \mathcal{B}_+^{*i}(z_2) : \frac{(q^{r^*} z_2/z_1; p^*)_\infty}{(q^{r^*+2} z_2/z_1; p^*)_\infty}, \\ : e^{\beta_2^i(z_1)} : \mathcal{B}_+^{*i}(z_2) &= : e^{\beta_2^i(z_1)} \mathcal{B}_+^{*i}(z_2) : \frac{(q^{r^*} z_2/z_1; p^*)_\infty}{(q^{r^*+2} z_2/z_1; p^*)_\infty}, \\ : e^{\beta_3^i(z_1)} : \mathcal{B}_+^{*i}(z_2) &= : e^{\beta_3^i(z_1)} \mathcal{B}_+^{*i}(z_2) : \frac{(q^{r^*} z_2/z_1; p^*)_\infty}{(q^{r^*-2} z_2/z_1; p^*)_\infty}, \\ : e^{\beta_4^i(z_1)} : \mathcal{B}_+^{*i}(z_2) &= : e^{\beta_4^i(z_1)} \mathcal{B}_+^{*i}(z_2) : \frac{(q^{r^*} z_2/z_1; p^*)_\infty}{(q^{r^*+2} z_2/z_1; p^*)_\infty}, \\ \mathcal{B}_-^i(z_1) : e^{\gamma^i(z_2)} : &= : \mathcal{B}_-^i(z_1) e^{\gamma^i(z_2)} : \frac{(q^{r+1} z_2/z_1; p)_\infty}{(q^{r-1} z_2/z_1; p)_\infty}, \\ \mathcal{B}_-^i(z_1) : e^{\beta_1^i(z_2)} : &= : \mathcal{B}_-^i(z_1) e^{\beta_1^i(z_2)} : \frac{(q^r z_2/z_1; p)_\infty}{(q^{r+2} z_2/z_1; p)_\infty}, \\ \mathcal{B}_-^i(z_1) : e^{\beta_2^i(z_2)} : &= : \mathcal{B}_-^i(z_1) e^{\beta_2^i(z_2)} : \frac{(q^r z_2/z_1; p)_\infty}{(q^{r+2} z_2/z_1; p)_\infty}, \\ \mathcal{B}_-^i(z_1) : e^{\beta_3^i(z_2)} : &= : \mathcal{B}_-^i(z_1) e^{\beta_3^i(z_2)} : \frac{(q^r z_2/z_1; p)_\infty}{(q^{r-2} z_2/z_1; p)_\infty}, \end{aligned}$$

$$\mathcal{B}_-^i(z_1) : e^{\beta_4^i(z_2)} : = : \mathcal{B}_-^i(z_1) e^{\beta_4^i(z_2)} : \frac{(q^r z_2/z_1; p)_\infty}{(q^{r-2} z_2/z_1; p)_\infty},$$

$$\begin{aligned} e^{b_+^i(z_1)} \mathcal{B}_+^{*i}(z_2) &= : e^{b_+^i(z_1)} \mathcal{B}_+^{*i}(z_2) : \frac{(q^{r^*} z_2/z_1; p^*)_\infty^2}{(q^{r^*+2} z_2/z_1; p^*)_\infty (q^{r^*-2} z_2/z_1; p^*)_\infty}, \\ \mathcal{B}_-^i(z_1) e^{b_-^i(z_2)} &= : \mathcal{B}_-^i(z_1) e^{b_-^i(z_2)} : \frac{(q^r z_2/z_1; p)_\infty^2}{(q^{r+2} z_2/z_1; p)_\infty (q^{r-2} z_2/z_1; p)_\infty}, \\ e^{a_+^i(z_1)} \mathcal{A}^{*i}(z_2) &= : e^{a_+^i(z_1)} \mathcal{A}^{*i}(z_2) : \frac{(q^{r^*+k+5} z_2/z_1; p^*)_\infty (q^{r^*-k-5} z_2/z_1; p^*)_\infty}{(q^{r^*+k+1} z_2/z_1; p^*)_\infty (q^{r^*-k-1} z_2/z_1; p^*)_\infty}, \\ e^{a_+^1(z_1)} \mathcal{A}^{*2}(z_2) &= : e^{a_+^1(z_1)} \mathcal{A}^{*2}(z_2) : \frac{(q^{r^*+k+2} z_2/z_1; p^*)_\infty (q^{r^*-k-2} z_2/z_1; p^*)_\infty}{(q^{r^*+k+4} z_2/z_1; p^*)_\infty (q^{r^*-k-4} z_2/z_1; p^*)_\infty}, \\ e^{a_+^2(z_1)} \mathcal{A}^{*1}(z_2) &= : e^{a_+^2(z_1)} \mathcal{A}^{*1}(z_2) : \frac{(q^{r^*+k+2} z_2/z_1; p^*)_\infty (q^{r^*-k-2} z_2/z_1; p^*)_\infty}{(q^{r^*+k+4} z_2/z_1; p^*)_\infty (q^{r^*-k-4} z_2/z_1; p^*)_\infty}, \\ \mathcal{A}^i(z_1) e^{a_-^i(z_2)} &= : \mathcal{A}^i(z_1) e^{a_-^i(z_2)} : \frac{(q^{r+k+5} z_2/z_1; p)_\infty (q^{r-k-5} z_2/z_1; p)_\infty}{(q^{r+k+1} z_2/z_1; p)_\infty (q^{r-k-1} z_2/z_1; p)_\infty}, \\ \mathcal{A}^1(z_1) e^{a_-^2(z_2)} &= : \mathcal{A}^1(z_1) e^{a_-^2(z_2)} : \frac{(q^{r+k+2} z_2/z_1; p)_\infty (q^{r-k-2} z_2/z_1; p)_\infty}{(q^{r+k+4} z_2/z_1; p)_\infty (q^{r-k-4} z_2/z_1; p)_\infty}, \\ \mathcal{A}^2(z_1) e^{a_-^1(z_2)} &= : \mathcal{A}^2(z_1) e^{a_-^1(z_2)} : \frac{(q^{r+k+2} z_2/z_1; p)_\infty (q^{r-k-2} z_2/z_1; p)_\infty}{(q^{r+k+4} z_2/z_1; p)_\infty (q^{r-k-4} z_2/z_1; p)_\infty}, \end{aligned}$$

$$\begin{aligned} \mathcal{B}_-^i(z_1) \mathcal{B}_+^{*i}(z_2) &= : \mathcal{B}_-^i(z_1) \mathcal{B}_+^{*i}(z_2) : \frac{(q^k z_2/z_1; q^{2k}, p^*)_\infty^2}{(q^{k+2} z_2/z_1; q^{2k}, p^*)_\infty (q^{k-2} z_2/z_1; q^{2k}, p^*)_\infty} \\ &\times \frac{(q^{k+2} z_2/z_1; q^{2k}, p)_\infty (q^{k-2} z_2/z_1; q^{2k}, p)_\infty}{(q^k z_2/z_1; q^{2k}, p)_\infty^2}, \\ \mathcal{A}^i(z_1) \mathcal{A}^{*i}(z_2) &= : \mathcal{A}^i(z_1) \mathcal{A}^{*i}(z_2) : \frac{(q^{2k+5} z_2/z_1; q^{2k}, p^*)_\infty (q^{-5} z_2/z_1; q^{2k}, p^*)_\infty}{(q^{2k+1} z_2/z_1; q^{2k}, p^*)_\infty (q^{-1} z_2/z_1; q^{2k}, p^*)_\infty} \\ &\times \frac{(q^{2k+1} z_2/z_1; q^{2k}, p)_\infty (q^{-1} z_2/z_1; q^{2k}, p)_\infty}{(q^{2k+5} z_2/z_1; q^{2k}, p)_\infty (q^{-5} z_2/z_1; q^{2k}, p)_\infty}, \\ \mathcal{A}^1(z_1) \mathcal{A}^{*2}(z_2) &= : \mathcal{A}^1(z_1) \mathcal{A}^{*2}(z_2) : \frac{(q^{2k+2} z_2/z_1; q^{2k}, p^*)_\infty (q^{-2} z_2/z_1; q^{2k}, p^*)_\infty}{(q^{2k+4} z_2/z_1; q^{2k}, p^*)_\infty (q^{-4} z_2/z_1; q^{2k}, p^*)_\infty} \\ &\times \frac{(q^{2k+4} z_2/z_1; q^{2k}, p)_\infty (q^{-4} z_2/z_1; q^{2k}, p)_\infty}{(q^{2k+2} z_2/z_1; q^{2k}, p)_\infty (q^{-2} z_2/z_1; q^{2k}, p)_\infty}, \\ \mathcal{A}^2(z_1) \mathcal{A}^{*1}(z_2) &= : \mathcal{A}^2(z_1) \mathcal{A}^{*1}(z_2) : \frac{(q^{2k+2} z_2/z_1; q^{2k}, p^*)_\infty (q^{-2} z_2/z_1; q^{2k}, p^*)_\infty}{(q^{2k+4} z_2/z_1; q^{2k}, p^*)_\infty (q^{-4} z_2/z_1; q^{2k}, p^*)_\infty} \\ &\times \frac{(q^{2k+4} z_2/z_1; q^{2k}, p)_\infty (q^{-4} z_2/z_1; q^{2k}, p)_\infty}{(q^{2k+2} z_2/z_1; q^{2k}, p)_\infty (q^{-2} z_2/z_1; q^{2k}, p)_\infty}. \end{aligned}$$

Here we have used the notation

$$(z; p_1, p_2)_\infty = \prod_{n_1, n_2=0}^{\infty} (1 - p_1^{n_1} p_2^{n_2} z).$$

References

- [1] O.Foda, K.Iohara, M.Jimbo, R.Kedem, T.Miwa, H.Yan: An elliptic quantum algebra for \widehat{sl}_2 , *Lett.Math.Phys.***32**, 259-268 (1994).
- [2] G.Felder: Elliptic quantum groups, Proceedings for International Conference of Mathematical Physics 1994, Paris, Cambridge-Hong Kong : International Press, 1995, pp.211-218.
- [3] C.Frønsdal : Quasi-Hopf deformation of quantum group, *Lett.Math.Phys.***40**, 117-134 (1997).
- [4] B.Enriquez and G.Felder : Elliptic quantum group $E_{\tau,\eta}(sl_2)$ and quasi-Hopf algebra, *Commun.Math.Phys.***195**, 651-689, (1998).
- [5] M.Jimbo, H.Konno, S.Odake and J.Shiraishi : Quasi-Hopf twistors for elliptic quantum groups, *Transformation Groups* **4**, 303-327, (1999).
- [6] V.Drinfeld : Quasi-Hopf algebras, *Leningrad Math.J.* **1**, 1419-1457, (1990).
- [7] H.Konno : An Elliptic Algebra $U_{q,p}(\widehat{sl}_2)$ and the fusion RSOS model, *Commun.Math.Phys.***195**, 373-403, (1998).
- [8] E.Date, M.Jimbo, A.Kuniba, T.Miwa and M.Okado : Exactly solvable SOS model: II. proof of the star triangle relation and combinatorial identities, *Adv.Stu.Pure.Math.***16**,17-122, (1998).
- [9] M.Jimbo, H.Konno, S.Odake and J.Shiraishi : Elliptic Algebra $U_{q,p}(\widehat{sl}_2)$: Drinfeld currents and vertex operators, *Commun.Math.Phys.***199**, 605-647, (1999).
- [10] T.Kojima and H.Konno : The Elliptic Algebra $U_{q,p}(\widehat{sl}_N)$ and the Drinfeld Realization of the Elliptic Quantum Group $\mathcal{B}_{q,\lambda}(\widehat{sl}_N)$, *Commun.Math.Phys.***237**, 405-447, (2003).
- [11] M.Wakimoto : Fock representation of the Affine Lie Algebra $A_1^{(1)}$, *Commun.Math.Phys.***104**, 605-609, (1986).
- [12] B.Feigin and E.Frenkel : Representation of Affine Kac-Moody algebra and bosonization, *Physics and Mathematics of Strings*, World Scientific, 1990, 271-316.

- [13] A.Matsuo : A q-deformation of Wakimoto modules, primary fields and screening operators, *Commun.Math.Phys.***160**,33-48,(1994).
- [14] J.Shiraishi : Free Boson Realization of $U_q(\widehat{sl}_2)$, *Phys.Lett.***A171**, 243-248, (1992).
- [15] H.Awata, S.Odake and J.Shiraishi : Free Boson Representation of $U_q(\widehat{sl}_3)$, *Lett.Math.Phys.***30**, 207-216, (1994).
- [16] W.Chang and X.Ding : On the vertex operators of the elliptic quantum algebra $U_{q,p}(\widehat{sl}_2)_k$ *J.Math.Phys.***49**, 043513, (2008).
- [17] T.Kojima and J.Shiraishi : The Integrals of Motion for the Deformed W -Algebra $W_{q,t}(\widehat{sl}_N)$.II.Proof of the Commutation Relations, *Commun.Math.Phys.***283**, 795-851, (2008).
- [18] T.Kojima and J.Shiraishi : A remark on the integrals of motion associated with level k realization of the elliptic algebra $U_{q,p}(\widehat{sl}_2)$, Proceedings for 10-th international conference on Geometry, Integrability and Quantization 2008, Bulgaria.
- [19] T.Kojima and J.Shiraishi : The integrals of motion for the elliptic deformation of the Virasoro and W_N algebra, Proceedings for 5-th World Congress for Nonlinear Analysts 2008, Florida, USA.